

A New Sequence Space Defined by a Sequence of Orlicz Functions over n -Normed Spaces

Kuldip RAJ, Sunil K. SHARMA

*School of Mathematics, Shri Mata Vaishno Devi University
Katra-182320, J&K, India
e-mail: kuldipraj68@gmail.com, sunilksharma42@yahoo.co.in*

(Received March 16, 2012)

Abstract

In this paper we introduce a new sequence space $BV_\sigma(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$ defined by a sequence of Orlicz functions $\mathcal{M} = (M_k)$ and study some topological properties of this sequence space.

Key words: paranorm space, invariant mean, orlicz function, Musielak–orlicz function, n -normed space, solid

2000 Mathematics Subject Classification: 40D05, 40A05

1 Introduction and Preliminaries

The concept of 2-normed spaces was initially developed by Gähler[2] in the mid of 1960's, while that of n -normed spaces one can see in Misiak[11]. Since then, many others have studied this concept and obtained various results, see Gunawan ([3], [4]), Gunawan and Mashadi [5] and many others. Let $n \in \mathbb{N}$ and X be a linear space of dimension d , where $d \geq n \geq 2$ over the field \mathbb{K} (\mathbb{K} is the field of real or complex numbers). A real valued function $\|\cdot, \dots, \cdot\|$ on X^n satisfying the following four conditions:

1. $\|x_1, x_2, \dots, x_n\| = 0$ if and only if x_1, x_2, \dots, x_n are linearly dependent in X ;
2. $\|x_1, x_2, \dots, x_n\|$ is invariant under permutation;
3. $\|\alpha x_1, x_2, \dots, x_n\| = |\alpha| \|x_1, x_2, \dots, x_n\|$ for any $\alpha \in \mathbb{K}$, and
4. $\|x + x', x_2, \dots, x_n\| \leq \|x, x_2, \dots, x_n\| + \|x', x_2, \dots, x_n\|$

is called a n -norm on X and the pair $(X, \|\cdot, \dots, \cdot\|)$ is called a n -normed space over the field \mathbb{K} .

For example, we may take $X = \mathbb{R}^n$ being equipped with the Euclidean n -norm, $\|x_1, x_2, \dots, x_n\|_E$ = the volume of the n -dimensional parallelepiped spanned by the vectors x_1, x_2, \dots, x_n which may be given explicitly by the formula

$$\|x_1, x_2, \dots, x_n\|_E = |\det(x_{ij})|,$$

where $x_i = (x_{i1}, x_{i2}, \dots, x_{in}) \in \mathbb{R}^n$ for each $i = 1, 2, \dots, n$ and $\|\cdot\|_E$ denotes the Euclidean norm. Let $(X, \|\cdot, \dots, \cdot\|)$ be an n -normed space of dimension $d \geq n \geq 2$ and $\{a_1, a_2, \dots, a_n\}$ be linearly independent set in X . Then the following function $\|\cdot, \dots, \cdot\|_\infty$ on X^{n-1} defined by

$$\|x_1, x_2, \dots, x_{n-1}\|_\infty = \max\{\|x_1, x_2, \dots, x_{n-1}, a_i\| : i = 1, 2, \dots, n\}$$

defines an $(n-1)$ -norm on X with respect to $\{a_1, a_2, \dots, a_n\}$.

A sequence (x_k) in a n -normed space $(X, \|\cdot, \dots, \cdot\|)$ is said to converge to some $L \in X$ if

$$\lim_{k \rightarrow \infty} \|x_k - L, z_1, \dots, z_{n-1}\| = 0 \quad \text{for every } z_1, \dots, z_{n-1} \in X.$$

A sequence (x_k) in a n -normed space $(X, \|\cdot, \dots, \cdot\|)$ is said to be Cauchy if

$$\lim_{k, p \rightarrow \infty} \|x_k - x_p, z_1, \dots, z_{n-1}\| = 0 \quad \text{for every } z_1, \dots, z_{n-1} \in X.$$

If every Cauchy sequence in X converges to some $L \in X$, then X is said to be complete with respect to the n -norm. A complete n -normed space is said to be a n -Banach space.

Let X be a linear metric space. A function $p: X \rightarrow \mathbb{R}$ is called paranorm, if

1. $p(x) \geq 0$, for all $x \in X$;
2. $p(-x) = p(x)$, for all $x \in X$;
3. $p(x + y) \leq p(x) + p(y)$, for all $x, y \in X$;
4. if (σ_n) is a sequence of scalars with $\sigma_n \rightarrow \sigma$ as $n \rightarrow \infty$ and (x_n) is a sequence of vectors with $p(x_n - x) \rightarrow 0$ as $n \rightarrow \infty$, then $p(\sigma_n x_n - \sigma x) \rightarrow 0$ as $n \rightarrow \infty$.

A paranorm p for which $p(x) = 0$ implies $x = 0$ is called total paranorm and the pair (X, p) is called a total paranormed space. It is well known that the metric of any linear metric space is given by some total paranorm (see [20], Theorem 10.4.2, P-183). For more details about sequence spaces see ([6], [12], [16], [18]).

Let l_∞ and c denotes the Banach spaces of bounded and convergent sequences $x = (x_k)_{k=1}^\infty$ respectively. Let σ be an injection of the set of positive integers \mathbb{N} into itself having no finite orbits and T be the operator defined on l_∞ by $T((x_n)_{n=1}^\infty) = (x_{\sigma(n)})_{n=1}^\infty$.

A positive linear functional φ , with $\|\varphi\| = 1$, is called a σ -mean or an invariant mean if $\varphi(x) = \varphi(Tx)$ for all $x \in l_\infty$.

A sequence $x = (x_k)$ is said to be σ -convergent, denoted by $x \in V_\sigma$, if $\varphi(x)$ takes the same value, called σ -lim x , for all σ -means φ . We have

$$V_\sigma = \left\{ x = (x_n) : \sum_{m=1}^{\infty} t_{m,n}(x) = L \text{ uniformly in } n, L = \sigma - \lim x \right\},$$

for $m \geq 0, n > 0$, where, $t_{m,n}(x) = \frac{x_n + x_{\sigma(n)} + \dots + x_{\sigma^m(n)}}{m+1}$, and $t_{-1,n} = 0$ (see schaefer [19]), where $\sigma^m(n)$ denotes the m^{th} iterate of σ at n . In particular, if σ is a translation, a σ -mean is often called a Banach limit and V_σ reduces to f , the set of almost convergent sequences (see Lorentz [8]). Subsequently invariant mean have been studied by Ahmad and Mursaleen [1] and many others.

A sequence space E is said to be solid(or normal) if $(x_k) \in E$ implies $(\alpha_k x_k) \in E$ for all sequences of scalars (α_k) with $|\alpha_k| \leq 1$ and for all $k \in \mathbb{N}$.

A sequence space E is said to be monotone if it contains the canonical preimages of all its step spaces.

An orlicz function M is a function, which is continuous, non-decreasing and convex with $M(0) = 0, M(x) > 0$ for $x > 0$ and $M(x) \rightarrow \infty$ as $x \rightarrow \infty$.

Lindenstrauss and Tzafriri [7] used the idea of Orlicz function to define the following sequence space. Let w be the space of all real or complex sequences $x = (x_k)$, then

$$\ell_M = \left\{ x \in w : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) < \infty \right\}$$

which is called as an Orlicz sequence space. The space ℓ_M is a Banach space with the norm

$$\|x\| = \inf \left\{ \rho > 0 : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) \leq 1 \right\}.$$

It is shown in [7] that every Orlicz sequence space ℓ_M contains a subspace isomorphic to $\ell_p (p \geq 1)$. The Δ_2 -condition is equivalent to $M(Lx) \leq kLM(x)$ for all values of $x \geq 0$, and for $L > 1$. An Orlicz function M can always be represented in the following integral form

$$M(x) = \int_0^x \eta(t) dt$$

where η is known as the kernel of M , is right differentiable for $t \geq 0, \eta(0) = 0, \eta(t) > 0, \eta$ is non-decreasing and $\eta(t) \rightarrow \infty$ as $t \rightarrow \infty$.

A sequence $\mathcal{M} = (M_k)$ of Orlicz functions is called a Musielak–Orlicz function see ([10], [14]). A sequence $\mathcal{N} = (N_k)$ defined by

$$N_k(v) = \sup\{|v|u - M_k(u) : u \geq 0\}, k = 1, 2, \dots$$

is called the complementary function of a Musielak–Orlicz function \mathcal{M} . For a given Musielak–Orlicz function \mathcal{M} , the Musielak–Orlicz sequence space $t_{\mathcal{M}}$ and

its subspace $h_{\mathcal{M}}$ are defined as follows

$$\begin{aligned} t_{\mathcal{M}} &= \{x \in w : I_{\mathcal{M}}(cx) < \infty \text{ for some } c > 0\}, \\ h_{\mathcal{M}} &= \{x \in w : I_{\mathcal{M}}(cx) < \infty \text{ for all } c > 0\}, \end{aligned}$$

where $I_{\mathcal{M}}$ is a convex modular defined by

$$I_{\mathcal{M}}(x) = \sum_{k=1}^{\infty} M_k(x_k), x = (x_k) \in t_{\mathcal{M}}.$$

We consider $t_{\mathcal{M}}$ equipped with the Luxemburg norm

$$\|x\| = \inf \left\{ k > 0 : I_{\mathcal{M}}\left(\frac{x}{k}\right) \leq 1 \right\}$$

or equipped with the Orlicz norm

$$\|x\|^0 = \inf \left\{ \frac{1}{k} (1 + I_{\mathcal{M}}(kx)) : k > 0 \right\}.$$

Mursaleen [13] defined the sequence space

$$BV_{\sigma} = \left\{ x \in l_{\infty} : \sum_m |\varphi_{m,n}(x)| < \infty, \text{ uniformly in } n \right\},$$

where $\varphi_{m,n}(x) = t_{m,n}(x) - t_{m-1,n}(x)$, assuming that $t_{m,n}(x) = 0$, for $m = -1$. Note that for any sequences $x = (x_k)$, $y = (y_k)$ and scalar λ we have

$$\varphi_{m,n}(x + y) = \varphi_{m,n}(x) + \varphi_{m,n}(y)$$

and

$$\varphi_{m,n}(\lambda x) = \lambda \varphi_{m,n}(x).$$

Let $\mathcal{M} = (M_k)$ be a Musielak–Orlicz function, $p = (p_m)$ be any sequence of strictly positive real numbers and $r \geq 0$ the sequence space $BV_{\sigma}(\mathcal{M}, p, r)$ defined by Raj, Sharma and Sharma [15].

Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions and $w(X)$ denotes X -valued sequence spaces. Let $p = (p_m)$ be a bounded sequence of positive real numbers and $u = (u_k)$ be any sequence of strictly positive real numbers. In the present paper we define the sequence space:

$$\begin{aligned} &BV_{\sigma}(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|) = \\ &\left\{ x = (x_k) \in w(X) : \sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} < \infty, \right. \\ &\left. \text{uniformly in } n \text{ and for some } \rho > 0 \right\}. \end{aligned}$$

For $\mathcal{M}(x) = x$, we get

$$BV_\sigma(u, p, r, \|\cdot, \dots, \cdot\|) = \\ \left\{ x = (x_k) \in w(X) : \sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} < \infty, \right. \\ \left. \text{uniformly in } n \text{ and for some } \rho > 0 \right\}.$$

For $p = p_m = 1$ for all m , we get

$$BV_\sigma(\mathcal{M}, u, r, \|\cdot, \dots, \cdot\|) = \\ \left\{ x = (x_k) \in w(X) : \sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right] < \infty, \right. \\ \left. \text{uniformly in } n \text{ and for some } \rho > 0 \right\}.$$

For $r = 0$, we get

$$BV_\sigma(\mathcal{M}, u, p, \|\cdot, \dots, \cdot\|) = \\ \left\{ x = (x_k) \in w(X) : \sum_{m=1}^{\infty} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} < \infty, \right. \\ \left. \text{uniformly in } n \text{ and for some } \rho > 0 \right\}.$$

For $\mathcal{M}(x) = x$ and $r = 0$, we get

$$BV_\sigma(u, p, \|\cdot, \dots, \cdot\|) = \\ \left\{ x = (x_k) \in w(X) : \sum_{m=1}^{\infty} \left[\sup_{k \geq 0} u_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} < \infty, \right. \\ \left. \text{uniformly in } n \text{ and for some } \rho > 0 \right\}.$$

For $p = p_m = 1$ for all m and $r = 0$, we get

$$BV_\sigma(\mathcal{M}, u, \|\cdot, \dots, \cdot\|) = \\ \left\{ x = (x_k) \in w(X) : \sum_{m=1}^{\infty} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right] < \infty, \right. \\ \left. \text{uniformly in } n \text{ and for some } \rho > 0 \right\}.$$

For $\mathcal{M}(x) = x$, $p = p_m = 1$ for all m and $r = 0$, we get,

$$BV_\sigma(u, \|\cdot, \dots, \cdot\|) = \\ \left\{ x = (x_k) \in w(X) : \sum_{m=1}^{\infty} \left[\sup_{k \geq 0} u_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right] < \infty, \right. \\ \left. \text{uniformly in } n \text{ and for some } \rho > 0 \right\}.$$

If we take $u = u_k = 1$ for all k we get,

$$BV_\sigma(\mathcal{M}, p, r, \|\cdot, \dots, \cdot\|) = \left\{ x = (x_k) \in w(X) : \sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho} \right\|, z_1, \dots, z_{n-1} \right) \right]^{p_m} < \infty, \right. \\ \left. \text{uniformly in } n \text{ and for some } \rho > 0 \right\}.$$

The following inequality will be used throughout the paper. If $0 \leq p_k \leq \sup p_k = H$, $K = \max(1, 2^{H-1})$ then

$$|a_k + b_k|^{p_k} \leq K \{|a_k|^{p_k} + |b_k|^{p_k}\} \quad (1)$$

for all k and $a_k, b_k \in \mathbb{C}$. Also $|a|^{p_k} \leq \max(1, |a|^H)$ for all $a \in \mathbb{C}$.

The aim of this paper is to examine some topological properties and inclusion relations between above defined sequence spaces.

2 Some properties of sequence space $BV_\sigma(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$

Theorem 2.1 *The sequence space $BV_\sigma(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$ is a linear space over the field of complex numbers \mathbb{C} .*

Proof Let $x, y \in BV_\sigma(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$ and $\alpha, \beta \in \mathbb{C}$. Then there exist positive numbers ρ_1 and ρ_2 such that

$$\sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho_1} \right\|, z_1, \dots, z_{n-1} \right) \right]^{p_m} < \infty, \\ \text{uniformly in } n \text{ and for some } \rho_1 > 0$$

and

$$\sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(y)}{\rho_2} \right\|, z_1, \dots, z_{n-1} \right) \right]^{p_m} < \infty, \\ \text{uniformly in } n \text{ and for some } \rho_2 > 0.$$

Define

$$\rho_3 = \max(2|\alpha|\rho_1, 2|\beta|\rho_2).$$

Since $\mathcal{M} = (M_k)$ is non-decreasing, convex and so by using inequality (1.1), we have

$$\begin{aligned}
& \sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(\alpha x + \beta y)}{\rho_3}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \\
& \leq \sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\alpha \varphi_{m,n}(x)}{\rho_3}, z_1, \dots, z_{n-1} \right\| \right. \right. \\
& \quad \left. \left. + \left\| \frac{\beta \varphi_{m,n}(y)}{\rho_3}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \\
& \leq K \sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho_1}, z_1, \dots, z_{n-1} \right\| \right) \right] \\
& \quad + K \sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(y)}{\rho_2}, z_1, \dots, z_{n-1} \right\| \right) \right] \\
& < \infty, \text{ uniformly in } n.
\end{aligned}$$

This proves that $BV_{\sigma}(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$ is a linear space over the field \mathbb{C} of complex numbers. \square

Theorem 2.2 *Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions, $p = (p_m)$ be a bounded sequence of positive real numbers and $u = (u_k)$ be a sequence of strictly positive real numbers, the space $BV_{\sigma}(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$ is a paranormed space with the paranorm defined by*

$$g(x) = \inf_{n \geq 1} \left\{ \rho^{\frac{pn}{H}} : \left(\sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \right)^{\frac{1}{H}} \leq 1, \right. \\
\left. \text{uniformly in } n \right\},$$

where $H = \max(1, \sup p_m)$.

Proof It is clear that $g(x) = g(-x)$. Since $M(0) = 0$, we get $g(0) = 0$. By using Theorem 2.1, for $\alpha = \beta = 1$, we get

$$g(x + y) \leq g(x) + g(y).$$

For the continuity of scalar multiplication, let $\lambda \neq 0$ be any complex numbers, then by definition, we have

$$g(\lambda x) = \inf_{n \geq 1} \left\{ \rho^{\frac{pn}{H}} : \left(\sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(\lambda x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \right)^{\frac{1}{H}} \leq 1, \right. \\
\left. \text{uniformly in } n \right\}.$$

$$g(\lambda x) = \inf_{n \geq 1} \left\{ (|\lambda|s)^{\frac{pn}{H}} : \left(\sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(\lambda x)}{s|\lambda|}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \right)^{\frac{1}{H}} \leq 1, \right. \\ \left. \text{uniformly in } n \right\},$$

where $s = \frac{\rho}{|\lambda|}$. Since $|\lambda|^{p_m} \leq \max(1, |\lambda|^q)$, we have

$$g(\lambda x) \leq \max(1, |\lambda|^q) \inf_{n \geq 1} \left\{ s^{\frac{pn}{H}} : \left(\sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}}{s}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \right)^{\frac{1}{H}} \leq 1, \right. \\ \left. \text{uniformly in } n \right\} = \max(1, |\lambda|^q) g(x),$$

and therefore $g(\lambda x)$ converges to zero in $BV_{\sigma}(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$. Now let x be fixed element in $BV_{\sigma}(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$, there exist $\rho > 0$ such that

$$g(x) = \inf_{n \geq 1} \left\{ \rho^{\frac{pn}{H}} : \left(\sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \right)^{\frac{1}{H}} \leq 1, \right. \\ \left. \text{uniformly in } n \right\}.$$

Now

$$g(\lambda x) = \inf_{n \geq 1} \left\{ \rho^{\frac{pn}{H}} : \left(\sum_{m=1}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(\lambda x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \right)^{\frac{1}{H}} \leq 1, \right. \\ \left. \text{uniformly in } n \right\} \rightarrow 0 \text{ as } \lambda \rightarrow 0.$$

This completes the proof. \square

Theorem 2.3 Suppose that $0 < p_m \leq q_m < \infty$, for each $m \in \mathbb{N}$ and $r \geq 0$. Then

- (i) $BV_{\sigma}(\mathcal{M}, u, p, \|\cdot, \dots, \cdot\|) \subseteq BV_{\sigma}(\mathcal{M}, u, q, \|\cdot, \dots, \cdot\|)$,
- (ii) $BV_{\sigma}(\mathcal{M}, u, \|\cdot, \dots, \cdot\|) \subseteq BV_{\sigma}(\mathcal{M}, u, r, \|\cdot, \dots, \cdot\|)$.

Proof (i) Suppose that $x \in BV_{\sigma}(\mathcal{M}, u, p, \|\cdot, \dots, \cdot\|)$. This implies that

$$\left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \leq 1$$

for sufficiently large value of m , say $m \geq m_0$ for some fixed $m_0 \in \mathbb{N}$. Since $\mathcal{M} = (M_k)$ is non-decreasing, we have

$$\begin{aligned} & \sum_{m=m_0}^{\infty} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{q_m} \\ & \leq \sum_{m=m_0}^{\infty} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z-1, \dots, z_{n-1} \right\| \right) \right]^{p_m} < \infty. \end{aligned}$$

Hence $x \in BV_{\sigma}(\mathcal{M}, u, q, \|\cdot, \dots, \cdot\|)$.

(ii) Suppose that $x \in BV_{\sigma}(\mathcal{M}, u, \|\cdot, \dots, \cdot\|)$. This implies that

$$\left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right] \leq 1,$$

for sufficiently large value of m , say $m = m_0$ for fixed $m_0 \in \mathbb{N}$. Since $\mathcal{M} = (M_k)$ is non-decreasing, we have

$$\begin{aligned} & \sum_{m=m_0}^{\infty} \frac{1}{m^r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right] \\ & \leq \sum_{m=m_0}^{\infty} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right] < \infty. \end{aligned}$$

Hence $x \in BV_{\sigma}(\mathcal{M}, u, r, \|\cdot, \dots, \cdot\|)$. \square

Corollary 2.1 (i) If $0 < p_m \leq 1$ for each m , then

$$BV_{\sigma}(\mathcal{M}, u, p, \|\cdot, \dots, \cdot\|) \subseteq BV_{\sigma}(\mathcal{M}, u, \|\cdot, \dots, \cdot\|).$$

(ii) If $p_m \geq 1$ for all m , then

$$BV_{\sigma}(\mathcal{M}, u, \|\cdot, \dots, \cdot\|) \subseteq BV_{\sigma}(\mathcal{M}, u, p, \|\cdot, \dots, \cdot\|).$$

Proof It follows from the above Theorem. \square

Theorem 2.4 The sequence space $BV_{\sigma}(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$ is solid.

Proof Let $x \in BV_{\sigma}(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$. This implies that

$$\sum_{m=1}^{\infty} m^{-r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} < \infty.$$

Let (α_m) be the sequence of scalars such that $|\alpha_m| \leq 1$ for all $m \in \mathbb{N}$. Then the result follows from the following inequality

$$\begin{aligned} & \sum_{m=1}^{\infty} m^{-r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\alpha_m \varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \\ & \leq \sum_{m=1}^{\infty} m^{-r} \left[\sup_{k \geq 0} u_k M_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} < \infty. \end{aligned}$$

Hence $\alpha x \in BV_\sigma(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$ for all sequences of scalars (α_m) with $|\alpha_m| \leq 1$ for all $m \in \mathbb{N}$ whenever $x \in BV_\sigma(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$. \square

Corollary 2.2 *The sequence space $BV_\sigma(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$ is monotone.*

Proof It follows from the above Theorem. \square

Theorem 2.5 *Let $\mathcal{M} = (M_k)$, $\mathcal{M}' = (M'_k)$, $\mathcal{M}'' = (M''_k)$ are sequences of Orlicz functions satisfying Δ_2 -condition and $r, r_1, r_2 \geq 0$. Then we have*

(i) *If $r > 1$ then $BV_\sigma(\mathcal{M}', u, p, r, \|\cdot, \dots, \cdot\|) \subseteq BV_\sigma(\mathcal{M} \circ \mathcal{M}', u, p, r, \|\cdot, \dots, \cdot\|)$,*

(ii) *$BV_\sigma(\mathcal{M}', u, p, r, \|\cdot, \dots, \cdot\|) \cap BV_\sigma(\mathcal{M}'', u, p, r, \|\cdot, \dots, \cdot\|)$*

$$\subseteq BV_\sigma(\mathcal{M}' + \mathcal{M}'', u, p, r, \|\cdot, \dots, \cdot\|),$$

(iii) *If $r_1 \leq r_2$ then $BV_\sigma(\mathcal{M}, u, p, r_1, \|\cdot, \dots, \cdot\|) \subseteq BV_\sigma(\mathcal{M}, u, p, r_2, \|\cdot, \dots, \cdot\|)$.*

Proof (i) Since $\mathcal{M}' = (M'_k)$ is continuous at origin from right for all k , for $\epsilon > 0$ there exists $0 < \delta < 1$ such that $0 \leq C \leq \delta$ implies $M'_k(C) < \epsilon$. If we define

$$I_1 = \left\{ m \in \mathbb{N} : \sup_{k \geq 0} u_k M'_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \leq \delta, \text{ for some } \rho > 0 \right\},$$

$$I_2 = \left\{ m \in \mathbb{N} : \sup_{k \geq 0} u_k M'_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) > \delta, \text{ for some } \rho > 0 \right\},$$

when $\sup_{k \geq 0} u_k M'_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) > \delta$, we get

$$\begin{aligned} & \sup_{k \geq 0} u_k M_k \left(\sup_{k \geq 0} M'_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right) \\ & \leq \left\{ 2 \sup_{k \geq 0} u_k M_k(1) / \delta \right\} \sup_{k \geq 0} u_k M'_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right). \end{aligned}$$

Hence for $x \in BV_\sigma(\mathcal{M}', u, p, r, \|\cdot, \dots, \cdot\|)$ and $r > 1$, we have

$$\begin{aligned} & \sum_{m=1}^{\infty} m^{-r} \left[\sup_{k \geq 0} u_k (M_k \circ M'_k) \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \\ & = \sum_{m \in I_1} m^{-r} \left[\sup_{k \geq 0} u_k (M_k \circ M'_k) \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \\ & + \sum_{m \in I_2} m^{-r} \left[\sup_{k \geq 0} u_k (M_k \circ M'_k) \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \\ & \leq \sum_{m \in I_1} m^{-r} [\epsilon]^{p_m} \\ & + \sum_{m \in I_2} m^{-r} \left[\left\{ \sup_{k \geq 0} 2u_k M_k(1) / \delta \right\} \sup_{k \geq 0} u_k M'_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \\ & \leq \max(\epsilon^h, \epsilon^H) \sum_{m=1}^{\infty} m^{-r} + \max \left(\left\{ \frac{2M_k(1)}{\delta} \right\}^h, \left\{ \frac{2M_k(1)}{\delta} \right\}^H \right), \end{aligned}$$

where $0 < h = \inf p_m \leq p_m \leq H = \sup p_m < \infty$.

(ii) The proof follows from the following inequality

$$\begin{aligned} & m^{-r} \left[\sup_{k \geq 0} u_k (M'_k + M''_k) \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \\ & \leq K m^{-r} \left[u_k M'_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m} \\ & \quad + K m^{-r} \left[u_k M''_k \left(\left\| \frac{\varphi_{m,n}(x)}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_m}. \end{aligned}$$

(iii) The proof is straight forward. \square

Corollary 2.3 Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions satisfying Δ_2 -condition. Then we have

- (i) If $r > 1$, then $BV_\sigma(u, p, r, \|\cdot, \dots, \cdot\|) \subset BV_\sigma(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$,
- (ii) $BV_\sigma(\mathcal{M}, u, p, \|\cdot, \dots, \cdot\|) \subseteq BV_\sigma(\mathcal{M}, u, p, r, \|\cdot, \dots, \cdot\|)$,
- (iii) $BV_\sigma(u, p, \|\cdot, \dots, \cdot\|) \subseteq BV_\sigma(u, p, r, \|\cdot, \dots, \cdot\|)$,
- (iv) $BV_\sigma(\mathcal{M}) \subseteq BV_\sigma(\mathcal{M}, u, r, \|\cdot, \dots, \cdot\|)$.

Proof The proof follows from the above theorem. \square

Acknowledgement The authors thank the referee for his valuable suggestions that improved the presentation of the paper.

References

- [1] Ahmad, Z. U., Mursaleen, M.: *An application of Banach limits*. Proc. Amer. Math. Soc., **103** (1983), 244–246.
- [2] Gähler S.: *Linear 2-normierte Rume*. Math. Nachr. **28** (1965), 1–43.
- [3] Gunawan H.: *On n -inner product, n -norms, and the Cauchy–Schwartz Inequality*. Scientiae Mathematicae Japonicae **5** (2001), 47–543.
- [4] Gunawan H.: *The space of p -summable sequence and its natural n -norm*. Bull. Aust. Math. Soc. **64** (2001), 137–147.
- [5] Gunawan H., Mashadi M.: *On n -normed spaces*. Int. J. Math. Math. Sci. **27** (2001), 631–639.
- [6] Khan, V. A.: *On a new sequence space defined by Orlicz functions*. Commun. Fac. Sci. Univ. Ank. A₁ **57** (2008), 25–33.
- [7] Lindenstrauss, J., Tzafriri, L.: *On Orlicz seequence spaces*. Israel J. Math. **10** (1971), 379–390.
- [8] Lorentz G. G.: *A Contribution to the theory of divergent series*. Acta Math. **80** (1948), 167–190.
- [9] Maddox, I. J.: *Some properties of paranormed sequence spaces*. J. London Math. Soc. **1**, (1989), 316–322.
- [10] Maligranda, L.: *Orlicz spaces & interpolation*. Seminars in mathematics **5** Polish Academy of Sciences, 1989.
- [11] Misiak A.: *n -inner product spaces*. Math. Nachr. **140**, (1989), 299–319.

- [12] Murasaleen, M.: *Matrix transformation between some new sequence spaces*. Houston J. Math. **9** (1983), 505–509.
- [13] Murasaleen, M.: *On some new invariant matrix methods of summability*. Quart. J. Math. Oxford **34**, (1983), 77–86.
- [14] Musielak, J.: Orlicz Spaces. Lectures Notes in Math. **1034** Springer, 1983.
- [15] Raj K., Sharma, A. K., Sharma S. K.: *A Sequence space defined by Musielak-Orlicz functions*. Int. J. Pure Appl. Math. **67** (2011), 475–484.
- [16] Raj K., Sharma, S. K., Sharma A. K.: *Some difference sequence spaces in n -normed spaces defined by Musielak-Orlicz function*. Armenian J. Math. **3** (2010), 127–141.
- [17] Raj K., Sharma, S. K.: *Some sequence spaces in 2-normed spaces defined by Musielak-Orlicz functions*. Acta Univ. Sapientiae Math. **3** (2011), 97–109.
- [18] Rao, M. M, Ren, Z. D: Theory of Orlicz Spaces. *Marcel Dekker Inc.* New york, Basel, Hongkong, 1991.
- [19] Schafer, P.: *Infinite matrices and invariant means*. Proc. Amer. Math. Soc. **36** (1972), 104–110.
- [20] Wilansky, A.: Summability through Functional Analysis. *North-Holland Math. Stud.*, 1984.