

Stability and Boundedness of Solutions of a Certain System of Third-order Nonlinear Delay Differential Equations

M. O. OMEIKE

*Department of Mathematics, Federal University of Agriculture
Abeokuta, Nigeria
e-mail: moomeike@yahoo.com*

(Received July 26, 2014)

Abstract

In this paper a number of known results on the stability and boundedness of solutions of some scalar third-order nonlinear delay differential equations are extended to some vector third-order nonlinear delay differential equations.

Key words: Lyapunov functional, third-order vector delay differential equation, boundedness, stability.

2010 Mathematics Subject Classification: 34K20

1 Introduction

The delay differential equation considered here is of the form

$$\ddot{X} + A\ddot{X} + B\dot{X} + H(X(t - r(t))) = P(t), \quad (1.1)$$

in which $X \in \mathbb{R}^n$, $P: \mathbb{R} \rightarrow \mathbb{R}^n$, A and B are real $n \times n$ constant matrices, $0 \leq r(t) \leq \gamma$, γ is a positive constant which will be determined later, and the dots indicate differentiation with respect to t . The equation is the vector version for the system of real third-order delay differential equations

$$\ddot{x}_i + \sum_{k=1}^n a_{ik} \ddot{x}_k + \sum_{k=1}^n b_{ik} \dot{x}_k + h_i(x_1(t - r(t)), x_2(t - r(t)), \dots, x_n(t - r(t))) = p_i(t),$$

$i = 1, 2, \dots, n$, in which a_{ik}, b_{ik} are constants. It will be assumed as basic throughout what follows that $H \in \mathcal{C}'(\mathbb{R}^n)$ and $P \in \mathcal{C}(\mathbb{R})$ are such that solutions of (1.1) exist corresponding to any pre-assigned initial conditions. Here, $\mathcal{C}'(\mathbb{R}^n)$

is the family of all functions once continuously differentiable on \mathbb{R}^n and $\mathcal{C}(\mathbb{R})$ is the family of all functions continuous on \mathbb{R} .

The study of (1.1) is concerned primarily with the problems of stability and boundedness of solutions of (1.1).

In the case $n = 1$, these problems have been investigated (see [3, 4, 5, 6, 10, 11, 14]) for a general scalar delay differential equation of the form

$$\ddot{x} + a\dot{x} + bx + h(x(t - r(t))) = p(t)$$

(a, b constants). Their investigation shows that the stability and the ultimate boundedness of solutions can be established if $h'(x)$ is bounded and if $a, b, h(x)$ satisfy some suitable generalization of the Routh–Hurwitz conditions

$$a > 0, \quad b > 0, \quad ab - c > 0$$

for the asymptotic stability of the solution $x = 0$ of the linear system

$$\ddot{x} + a\dot{x} + bx + cx = 0$$

with constant coefficients. The object of the present paper is to provide analogous results for n -dimensional equation (1.1) following the arguments used in some of the papers mentioned above.

Notation and definitions

Given any X, Y in \mathbb{R}^n the symbol $\langle X, Y \rangle$ will be used to denote the usual scalar product in \mathbb{R}^n , that is $\langle X, Y \rangle = \sum_{i=1}^n x_i y_i$; thus $\|X\|^2 = \langle X, X \rangle$. The matrix A is said to be positive definite when $\langle AX, X \rangle > 0$ for all nonzero X in \mathbb{R}^n .

The following notations (see [5, 15]) will be useful in subsequent sections. For $x \in \mathbb{R}^n$, $|x|$ is the norm of x . For a given $r > 0$, $t_1 \in \mathbb{R}$,

$$C(t_1) = \{\phi: [t_1 - r, t_1] \rightarrow \mathbb{R}^n / \phi \text{ is continuous}\}.$$

In particular, $C = C(0)$ denotes the space of continuous functions mapping the interval $[-r, 0]$ into \mathbb{R}^n and for $\phi \in C$, $\|\phi\| = \sup_{-r \leq \theta \leq 0} |\phi(\theta)|$. $C_{\mathbf{H}}$ will denote the set of ϕ such that $\|\phi\| \leq \mathbf{H}$. For any continuous function $x(u)$ defined on $-h \leq u < A$, $A > 0$, and any fixed t , $0 \leq t < A$, the symbol x_t will denote the restriction of $x(u)$ to the interval $[t - r, t]$, that is, x_t is an element of C defined by $x_t(\theta) = x(t + \theta)$, $-r \leq \theta \leq 0$.

2 Some preliminary results

In this section, we shall state the algebraic results required in the proofs of our main results. The proofs are not given since they are found in [1, 2, 7, 8, 9, 13].

Lemma 2.1 [1, 2, 7, 8, 9, 13] *Let D be a real symmetric positive definite $n \times n$ matrix, then for any X in \mathbb{R}^n , we have*

$$\delta_d \|X\|^2 \leq \langle DX, X \rangle \leq \Delta_d \|X\|^2$$

where δ_d, Δ_d are the least and the greatest eigenvalues of D , respectively.

Lemma 2.2 [1, 2, 7, 8, 9, 13] *Let Q, D be any two real $n \times n$ commuting symmetric matrices. Then*

(i) *the eigenvalues $\lambda_i(QD)$ ($i = 1, 2, \dots, n$) of the product matrix QD are all real and satisfy*

$$\min_{1 \leq j, k \leq n} \lambda_j(Q)\lambda_k(D) \leq \lambda_i(QD) \leq \max_{1 \leq j, k \leq n} \lambda_j(Q)\lambda_k(D);$$

(ii) *the eigenvalues $\lambda_i(Q + D)$ ($i = 1, 2, \dots, n$) of the sum of matrices Q and D are real and satisfy*

$$\left\{ \min_{1 \leq j \leq n} \lambda_j(Q) + \min_{1 \leq k \leq n} \lambda_k(D) \right\} \leq \lambda_i(Q + D) \leq \left\{ \max_{1 \leq j \leq n} \lambda_j(Q) + \max_{1 \leq k \leq n} \lambda_k(D) \right\}.$$

Lemma 2.3 [1, 2, 7, 8, 9, 13] *Let $H(X)$ be a continuous vector function and that $H(0) = 0$ then*

$$\frac{d}{dt} \int_0^1 \langle H(\sigma X), X \rangle d\sigma = \langle H(X), \dot{X} \rangle.$$

Lemma 2.4 *Let $H(X)$ be a continuous vector function and that $H(0) = 0$ then*

$$\delta_h \|X\|^2 \leq 2 \int_0^1 \langle H(\sigma X), X \rangle d\sigma \leq \Delta_h \|X\|^2$$

where δ_h, Δ_h are the least and the greatest eigenvalues of $J_h(X)$ (Jacobian matrix of H), respectively.

3 Stability

First, we will give the stability criteria for the general autonomous delay differential system. We consider

$$x' = f(x_t), \quad x_t(\theta) = x(t + \theta), \quad -r \leq \theta \leq 0, \quad t \geq 0, \tag{3.1}$$

where $f: C_{\mathbf{H}} \rightarrow \mathbb{R}^n$ is a continuous mapping, $f(0) = 0$,

$$C_{\mathbf{H}} := \{ \phi \in C([-r, 0], \mathbb{R}^n) : \|\phi\| \leq \mathbf{H} \}$$

and for $\mathbf{H}_1 < \mathbf{H}$, there exists $L > 0$, with $|f(\phi)| \leq L$ when $\|\phi\| \leq \mathbf{H}_1$. Here, $C([-r, 0], \mathbb{R}^n)$ is the family of all vector functions mapping $[-r, 0]$ into \mathbb{R}^n .

Definition 3.1 [3, 5, 6, 11, 12] *An element $\psi \in C$ is in the ω -limit set of ϕ , if $x(t, 0, \phi)$ is defined on $[0, \infty)$ and there is a sequence $\{t_n\}$, $t_n \rightarrow \infty$, as $n \rightarrow \infty$, with $\|x_{t_n}(\phi) - \psi\| \rightarrow 0$ as $n \rightarrow \infty$ where $x_{t_n}(\phi) = x(t_n + \theta, 0, \phi)$ for $-r \leq \theta \leq 0$. $x(t; 0, \phi)$ is a motion of a system at $t \in \mathbb{R}$ if and only if $x(0) = \phi$.*

Definition 3.2 [3, 5, 6, 11, 12] *A set $Q \subset C_{\mathbf{H}}$ is an invariant set if for any $\phi \in Q$, the solution of (3.1), $x(t, 0, \phi)$, is defined on $[0, \infty)$, and $x_t(\phi) \in Q$ for $t \in [0, \infty)$.*

Lemma 3.3 (see [3, 5, 6, 11, 12]) *If $\phi \in C_{\mathbf{H}}$ is such that the solution $x_t(\theta)$ of (3.1) with $x_0(\phi) = \phi$ is defined on $[0, \infty)$ and $\|x_t(\phi)\| \leq \mathbf{H}_1 < \mathbf{H}$ for $t \in [0, \infty)$, then $\Omega(\phi)$ (the ω -limit set of ϕ) is a nonempty, compact, invariant set and*

$$\text{dist}(x_t(\phi), \Omega(\phi)) \rightarrow 0, \quad \text{as } t \rightarrow \infty.$$

Lemma 3.4 (see [3, 5, 6, 11, 12]) *Let $V(\phi): C_{\mathbf{H}} \rightarrow \mathbb{R}$ be a continuous functional satisfying a local Lipschitz condition. $V(0) = 0$ and such that*

- (i) $W_1(\|\phi(0)\|) \leq V(\phi) \leq W_2(\|\phi(0)\|)$ where $W_1(r), W_2(r)$ are wedges.
- (ii) $V'_{(3.1)}(\phi) \leq 0$, for $\phi \in C_{\mathbf{H}}$.

Then the zero solution of (3.1) is uniformly stable. If we define

$$Z = \{\phi \in C_{\mathbf{H}}: V'_{(3.1)}(\phi) = 0\},$$

then the zero solution of (3.1) is asymptotically stable, provided that the largest invariant set in Z is $Q = \{0\}$.

Before we state our result in this section, we write equation (1.1) with $P \equiv 0$ as

$$\begin{aligned} \dot{X} &= Y \\ \dot{Y} &= Z \\ \dot{Z} &= -AZ - BY - H(X) + \int_{t-r(t)}^t J_h(X)Y \, ds. \end{aligned} \tag{3.2}$$

We shall constantly refer to (3.2) subsequently in our discussion.

The following will be our main stability result for (3.2).

Theorem 3.5 *Consider (3.2), let $H(0) = 0$ and suppose that*

- (i) $0 \leq r(t) \leq \gamma$ ($\gamma > 0$), $r'(t) \leq \xi$, and $0 < \xi < 1$;
- (ii) the matrices A, B and $J_h(X)$ (Jacobian matrix of $H(X)$) are symmetric and positive definite, and furthermore the eigenvalues $\lambda_i(A)$, $\lambda_i(B)$ and $\lambda_i(J_h(X))$ ($i = 1, 2, \dots, n$) of A, B and $J_h(X)$, respectively satisfy,

$$0 < \delta_a \leq \lambda_i(A) \leq \Delta_a \tag{3.3}$$

$$0 < \delta_b \leq \lambda_i(B) \leq \Delta_b \tag{3.4}$$

$$0 < \delta_h \leq \lambda_i(J_h(X)) \leq \Delta_h, \quad \text{for } X \in \mathbb{R}^n, \tag{3.5}$$

where $\delta_a, \delta_b, \delta_h, \Delta_a, \Delta_b, \Delta_h$ are finite constants;

- (iii) the matrices A, B and $J_h(X)$ commute pairwise.

Then the zero solution of (3.2) is asymptotically stable, provided

$$\gamma < \min \left\{ \frac{2(\beta\delta_a - 1)}{\beta\Delta_h}, \frac{2(\delta_b - \beta\Delta_h)(1 - \xi)}{\Delta_h(2 + \beta - \xi)} \right\}.$$

Proof Using the equivalent system form (3.2), our main tool is the following Lyapunov functional, $V_1(X_t, Y_t, Z_t)$ defined as

$$2V_1(X_t, Y_t, Z_t) = 2 \int_0^1 \langle H(\sigma X), X \rangle d\sigma + \langle AY, Y \rangle + \beta \langle Y, BY \rangle \\ + \beta \langle Z, Z \rangle + 2 \langle Y, Z \rangle + 2\beta \langle Y, H(X) \rangle + \mu \int_{-r(t)}^0 \int_{t+s}^t \langle Y(\theta), Y(\theta) \rangle d\theta ds, \quad (3.6)$$

where

$$\frac{1}{\delta_a} < \beta < \frac{\delta_b}{\Delta_h}.$$

Since

$$\mu \int_{-r(t)}^0 \int_{t+s}^t \langle Y(\theta), Y(\theta) \rangle d\theta ds$$

is non-negative, we have

$$2V_1(X_t, Y_t, Z_t) \geq 2 \int_0^1 \langle H(\sigma X), X \rangle d\sigma + \langle AY, Y \rangle + \beta \langle Y, BY \rangle \\ + \beta \langle Z, Z \rangle + 2 \langle Y, Z \rangle + 2\beta \langle Y, H(X) \rangle \\ = 2 \int_0^1 \langle H(\sigma X), X \rangle d\sigma - \beta \langle B^{-1}H(X), H(X) \rangle \\ + \beta \|B^{\frac{1}{2}}Y + B^{-\frac{1}{2}}H(X)\|^2 + \beta \|Z + \beta^{-1}Y\|^2 + \langle (A - \beta^{-1}I)Y, Y \rangle.$$

Using Lemmas 2.1, 2.2 and 2.4, and since $\beta \|B^{\frac{1}{2}}Y + B^{-\frac{1}{2}}H(X)\|^2 \geq 0$, we have

$$2V_1(X_t, Y_t, Z_t) \geq 2 \int_0^1 \int_0^1 \sigma \langle \{I - \beta B^{-1}J_h(\sigma X)\} J_h(\sigma \tau X) X, X \rangle d\sigma d\tau \\ + \beta \|Z + \beta^{-1}Y\|^2 + \langle (A - \beta^{-1}I)Y, Y \rangle \\ \geq (1 - \beta \delta_b^{-1} \Delta_h) \delta_h \|X\|^2 + \beta \|Z + \beta^{-1}Y\|^2 + (\delta_a - \beta^{-1}) \|Y\|^2.$$

Hence there is a constant $K > 0$ (small enough) such that

$$V_1(X_t, Y_t, Z_t) \geq K (\|X\|^2 + \|Y\|^2 + \|Z\|^2).$$

Next, our target is to show that $V_1(X_t, Y_t, Z_t)$ satisfies the second condition of Lemma 3.4. From (3.2), (3.6) and using Lemma 2.3, we have

$$\frac{d}{dt} V_1(X_t, Y_t, Z_t) = -\langle (\beta A - I)Z, Z \rangle - \langle \{B - \beta J_h(X)\} Y, Y \rangle \\ + \int_{t-r(t)}^t \langle Y, J_h(X)Y \rangle ds + \beta \int_{t-r(t)}^t \langle Z, J_h(X)Y \rangle ds \\ + \mu r(t) \langle Y, Y \rangle - \mu(1 - r'(t)) \int_{t-r(t)}^t \langle Y(\theta), Y(\theta) \rangle d\theta. \quad (3.7)$$

On using Lemmas 2.1 and 2.2, and the identity $2|\langle U, V \rangle| \leq \|U\|^2 + \|V\|^2$, we obtain,

$$\begin{aligned} \frac{d}{dt}V_1(X_t, Y_t, Z_t) &\leq - \left(\delta_b - \beta\Delta_h - \frac{1}{2}\Delta_h\gamma - \mu\gamma \right) \|Y\|^2 \\ &\quad - (\beta\delta_a - 1 - \frac{1}{2}\beta\gamma\Delta_h) \|Z\|^2 \\ &\quad + \left\{ \frac{1}{2}\beta\Delta_h + \frac{1}{2}\Delta_h - \mu(1 - \xi) \right\} \int_{t-r(t)}^t \langle Y(\theta), Y(\theta) \rangle d\theta. \end{aligned} \quad (3.8)$$

If we choose $\mu = \frac{(\beta+1)\Delta_h}{2(1-\xi)}$,

$$\begin{aligned} \frac{d}{dt}V_1(X_t, Y_t, Z_t) &\leq - \left\{ \delta_b - \beta\Delta_h - \frac{(2 + \beta - \xi)\Delta_h\gamma}{2(1 - \xi)} \right\} \|Y\|^2 \\ &\quad - \left(\beta\delta_a - 1 - \frac{1}{2}\beta\gamma\Delta_h \right) \|Z\|^2, \end{aligned}$$

and choosing

$$\gamma < \min \left\{ \frac{2(\beta\delta_a - 1)}{\beta\Delta_h}, \frac{2(\delta_b - \beta\Delta_h)}{\Delta_h(2 + \beta - \xi)} \right\},$$

there is a constant $D > 0$ such that

$$\frac{d}{dt}V_1(X_t, Y_t, Z_t) \leq -D (\|Y\|^2 + \|Z\|^2).$$

Hence the result follows. \square

Example 1 As a special case of system (1.1) (for $P(t) = 0$), let us take $n = 2$ that

$$X = \begin{pmatrix} x_1(t) \\ x_2(t) \end{pmatrix}, \quad A = \begin{pmatrix} 8 & 0 \\ 0 & 10 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix}$$

and

$$H(X(t - r(t))) = \begin{pmatrix} \tan^{-1} x_1(t - r(t)) + 2x_1(t - r(t)) \\ \tan^{-1} x_2(t - r(t)) + 2x_2(t - r(t)) \end{pmatrix}.$$

Thus,

$$H(X(t)) = \begin{pmatrix} \tan^{-1} x_1(t) + 2x_1(t) \\ \tan^{-1} x_2(t) + 2x_2(t) \end{pmatrix} \quad \text{and} \quad J_h(X) = \begin{pmatrix} 2 + \frac{1}{1+x_1^2} & 0 \\ 0 & 3 + \frac{1}{1+x_2^2} \end{pmatrix}.$$

If we take $r(t) = \frac{1}{22+t^2}$, then $0 \leq \frac{1}{22+t^2} < \gamma$, and that $r'(t) = \frac{-2t}{(22+t^2)^2} \leq \xi$, $0 < \xi < 1$. Clearly, A , B and $J_h(X)$ are symmetric and commute pairwise. That is,

$$\begin{aligned} AB &= \begin{pmatrix} 8 & 0 \\ 0 & 30 \end{pmatrix} = BA, \\ AJ_h(X) &= \begin{pmatrix} 16 + \frac{8}{1+x_1^2} & 0 \\ 0 & 30 + \frac{10}{1+x_2^2} \end{pmatrix} = J_h(X)A \end{aligned}$$

and

$$BJ_h(X) = \begin{pmatrix} 2 + \frac{1}{1+x_1^2} & 0 \\ 0 & 9 + \frac{3}{1+x_2^2} \end{pmatrix} = J_h(X)B.$$

Then, by easy calculation, we obtain eigenvalues of the matrices A, B and $J_h(X)$ as follows:

$$\lambda_1(A) = 8, \quad \lambda_2(A) = 10, \quad \lambda_1(B) = 1, \quad \lambda_2(B) = 3,$$

$$\lambda_1(J_h(X)) = 2 + \frac{1}{1+x_1^2}, \quad \lambda_2(J_h(X)) = 3 + \frac{1}{1+x_1^2}.$$

It is obvious that $\delta_a = 8, \Delta_a = 10, \delta_b = 1, \Delta_b = 3, \delta_h = 2, \Delta_h = 4$. If we choose $\beta = \frac{1}{6}$ and $\xi = \frac{1}{2}$, we must have that $\gamma < \min\{1, \frac{1}{20}\}$.

Thus, all the conditions of Theorem 3.5 are satisfied.

4 Boundedness

First, consider a system of delay differential equations

$$\dot{x} = F(t, x_t), \quad x_t(\theta) = x(t + \theta), \quad -r \leq \theta \leq 0, \quad t \geq 0, \quad (4.1)$$

where $F: \mathbb{R} \times C_{\mathbf{H}} \rightarrow \mathbb{R}^n$ is a continuous mapping and takes bounded set into bounded sets.

The following lemma is a well-known result obtained by Burton [5].

Lemma 4.1 [3, 5, 6, 11, 12] *Let $V(t, \phi): \mathbb{R} \times C_{\mathbf{H}} \rightarrow \mathbb{R}$ be continuous and locally Lipschitz in ϕ . If*

$$(i) \quad W(|x(t)|) \leq V(t, x_t) \leq W_1(|x(t)|) + W_2 \left(\int_{t-r(t)}^t W_3(|x(s)|) ds \right), \text{ and}$$

$$(ii) \quad \dot{V}_{(4.1)} \leq -W_3(|x(s)|) + M,$$

for some $M > 0$, where $W(r), W_i (i = 1, 2, 3)$ are wedges, then the solutions of (4.1) are uniformly bounded and uniformly ultimately bounded for bound \mathbf{B} .

To study the boundedness of solutions of (1.1) for which $P(t) \neq 0$, we would need to write Eq. (1.1) in the form

$$\begin{aligned} \dot{X} &= Y \\ \dot{Y} &= Z \\ \dot{Z} &= -AZ - BY - H(X) + \int_{t-r(t)}^t J_h(X)Y ds + P(t). \end{aligned} \quad (4.2)$$

Thus our main theorem in this section is stated with respect to Eq.(4.2) as follows:

Theorem 4.2 *If the conditions of Theorem 3.5 hold, and*

$$\|P(t)\| \leq m, \quad (4.3)$$

where m is a positive constant, then the solutions of equation (4.2) are uniformly bounded and uniformly ultimately bounded provided γ satisfies

$$\gamma < \min \left\{ \frac{2\delta_h(\delta_a\delta_b - \Delta_h)}{\Delta_h(\Delta_a\Delta_b - \Delta_h)}, \frac{2(\beta\delta_a - 1)}{\Delta_h(\beta + \Delta_a)}, \frac{2(\delta_b - \beta\Delta_h)(1 - \xi)}{\Delta_h \{(1 + \Delta_a^2)(1 - \xi) + 2 + \beta + \Delta_a^2 + (\Delta_a\Delta_b - \Delta_h)\}} \right\},$$

Proof Consider the function

$$V = V_1(X_t, Y_t, Z_t) + V_2(X_t, Y_t, Z_t), \quad (4.4)$$

where $V_1(X_t, Y_t, Z_t)$ is defined as (3.6) and $V_2(X_t, Y_t, Z_t)$ defined as

$$\begin{aligned} 2V_2(X_t, Y_t, Z_t) &= 2 \int_0^1 \langle AH(\sigma X), AX \rangle d\sigma + \langle B(AB - \Delta_h I)X, X \rangle \\ &\quad + 2\langle AY, H(X) \rangle + \langle \Delta_h Y, Y \rangle \\ &\quad + 2\langle (AB - \Delta_h I)X, Z + AY \rangle + \langle A(Z + AY), Z + AY \rangle. \end{aligned} \quad (4.5)$$

This we can rewrite

$$\begin{aligned} 2V_2(X_t, Y_t, Z_t) &= 2 \int_0^1 \langle AH(\sigma X), AX \rangle d\sigma - \langle A\Delta_h^{-1}H(X), AH(X) \rangle \\ &\quad + \langle \Delta_h Y + H(X), Y + A\Delta_h^{-1}H(X) \rangle + \langle \Delta_h A^{-1}(AB - \Delta_h I)X, X \rangle \\ &\quad + \langle (AB - \Delta_h I)X + A^2Y + AZ, A^{-1}(AB - \Delta_h I)X + AY + Z \rangle \\ &= 2 \int_0^1 \int_0^1 \sigma \langle \{I - \Delta_h^{-1}J_h(\sigma X)\} A^2 J(\tau\sigma X)X, X \rangle d\sigma d\tau \\ &\quad + \|A^{-\frac{1}{2}}(AB - \Delta_h I)X + A^{\frac{3}{2}}Y + A^{\frac{1}{2}}Z\|^2 \\ &\quad + \|\Delta_h^{\frac{1}{2}}(Y + \Delta_h^{-1}AH(X))\|^2 + \langle \Delta_h A^{-1}(AB - \Delta_h I)X, X \rangle. \end{aligned}$$

However,

$$2 \int_0^1 \int_0^1 \sigma \langle \{I - \Delta_h^{-1}J_h(\sigma X)\} A^2 J(\tau\sigma X)X, X \rangle d\sigma d\tau \geq 0$$

and $\langle \Delta_h A^{-1}(AB - \Delta_h I)X, X \rangle \geq \Delta_h \Delta_a^{-1}(\delta_a\delta_b - \Delta_h)\|X\|^2$. Thus,

$$\begin{aligned} 2V_2(X_t, Y_t, Z_t) &\geq \Delta_h \Delta_a^{-1}(\delta_a\delta_b - \Delta_h)\|X\|^2 + \|\Delta_h^{\frac{1}{2}}(Y + \Delta_h^{-1}AH(X))\|^2 \\ &\quad + \|A^{-\frac{1}{2}}(AB - \Delta_h I)X + A^{\frac{3}{2}}Y + A^{\frac{1}{2}}Z\|^2. \end{aligned}$$

It follows that $V_2(X_t, Y_t, Z_t)$ is positive definite.

From (3.6), (3.7), (3.8), (4.2) and Lemma 2.3, we find

$$\begin{aligned} \frac{d}{dt}V_1(X_t, Y_t, Z_t) &\leq - \left(\delta_b - \beta\Delta_h - \frac{1}{2}\Delta_h\gamma - \mu\gamma \right) \|Y\|^2 \\ &\quad - \left(\beta\delta_a - 1 - \frac{1}{2}\beta\gamma\Delta_h \right) \|Z\|^2 + \left\{ \frac{1}{2}\beta\Delta_h + \frac{1}{2}\Delta_h - \mu(1 - \xi) \right\} \\ &\quad \times \int_{t-r(t)}^t \langle Y(\theta), Y(\theta) \rangle d\theta + (\|Y\| + \beta\|Z\|)m. \end{aligned} \quad (4.6)$$

Also from (4.2), (4.5) and Lemma 2.3 we obtain,

$$\begin{aligned} \frac{d}{dt}V_2(X_t, Y_t, Z_t) &= -\langle (AB - \Delta_h I)X, H(X) \rangle - \langle A(\Delta_h I - J_h(X))Y, Y \rangle \\ &\quad + \int_{t-r(t)}^t \langle (AB - \Delta_h I)X, J_h(X)Y \rangle ds + \int_{t-r(t)}^t \langle AZ, J_h(X)Y \rangle ds \\ &\quad + \int_{t-r(t)}^t \langle A^2Y, J_h(X)Y \rangle ds + \langle P(t), (AB - \Delta_h I)X + A^2Y + AZ \rangle \end{aligned}$$

Also using Lemmas 2.1 and 2.2, the identity $2|\langle U, V \rangle| \leq (\|U\|^2 + \|V\|^2)$, and the fact that $\langle A(\Delta_h I - J_h(X))Y, Y \rangle \geq 0$ we find that

$$\begin{aligned} \frac{d}{dt}V_2(X_t, Y_t, Z_t) &\leq - \left\{ \delta_h(\delta_a\delta_b - \Delta_h) - \frac{1}{2}\Delta_h(\Delta_a\Delta_b - \Delta_h)r(t) \right\} \|X\|^2 \\ &\quad + \frac{1}{2}\Delta_a\Delta_h r(t)\|Z\|^2 + \frac{1}{2}\Delta_a^2\Delta_h r(t)\|Y\|^2 \\ &\quad + \frac{1}{2}\Delta_h \left\{ (\Delta_a\Delta_b - \Delta_h) + \Delta_a + \Delta_a^2 \right\} \int_{t-r(t)}^t \langle Y(s), Y(s) \rangle ds \\ &\quad + \left\{ (\Delta_a\Delta_b - \Delta_h)\|X\| + \Delta_a^2\|Y\| + \Delta_a\|Z\| \right\} m \end{aligned} \quad (4.7)$$

Therefore, from (4.4), (4.6) and (4.7), we obtain

$$\begin{aligned} \frac{d}{dt}V(X_t, Y_t, Z_t) &\leq - \left\{ \delta_h(\delta_a\delta_b - \Delta_h) - \frac{1}{2}\Delta_h\gamma(\Delta_a\Delta_b - \Delta_h) \right\} \|X\|^2 \\ &\quad - \left\{ \delta_b - \beta\Delta_h - \gamma \left(\frac{1}{2}\Delta_h + \frac{1}{2}\Delta_a^2\Delta_h + \mu \right) \right\} \|Y\|^2 \\ &\quad - \left\{ \beta\delta_a - 1 - \frac{1}{2}\gamma\Delta_h(\beta + \Delta_a) \right\} \|Z\|^2 \\ &\quad + \left\{ \frac{1}{2}\Delta_h (2 + \beta + \Delta_a^2 + (\Delta_a\Delta_b - \Delta_h)) - \mu(1 - \xi) \right\} \int_{t-r(t)}^t \langle Y(\theta), Y(\theta) \rangle d\theta \\ &\quad + \left\{ (\Delta_a\Delta_b - \Delta_h)\|X\| + (1 + \Delta_a^2)\|Y\| + (\beta + \Delta_a)\|Z\| \right\} m. \end{aligned}$$

If we choose

$$\mu = \frac{\Delta_h \{2 + \beta + \Delta_a^2 + (\Delta_a \Delta_b - \Delta_h)\}}{2(1 - \xi)},$$

we obtain

$$\begin{aligned} \frac{d}{dt}V(X_t, Y_t, Z_t) &\leq - \left\{ \delta_h(\delta_a \delta_b - \Delta_h) - \frac{1}{2} \Delta_h \gamma (\Delta_a \Delta_b - \Delta_h) \right\} \|X\|^2 \\ &\quad - \frac{1}{2(1 - \xi)} \{2(\delta_b - \beta \Delta_h)(1 - \xi) \\ &\quad - \gamma \Delta_h \{(1 + \Delta_a^2)(1 - \xi) + 2 + \beta + \Delta_a^2 + (\Delta_a \Delta_b - \Delta_h)\}\} \|Y\|^2 \\ &\quad - \left\{ \beta \delta_a - 1 - \frac{1}{2} \gamma \Delta_h (\beta + \Delta_a) \right\} \|Z\|^2 \\ &\quad + \{(\Delta_a \Delta_b - \Delta_h)\|X\| + (1 + \Delta_a^2)\|Y\| + (\beta + \Delta_a)\|Z\|\} m. \end{aligned}$$

When

$$\gamma < \min \left\{ \frac{2\delta_h(\delta_a \delta_b - \Delta_h)}{\Delta_h(\Delta_a \Delta_b - \Delta_h)}, \frac{2(\beta \delta_a - 1)}{\Delta_h(\beta + \Delta_a)}, \frac{2(\delta_b - \beta \Delta_h)(1 - \xi)}{\Delta_h \{(1 + \Delta_a^2)(1 - \xi) + 2 + \beta + \Delta_a^2 + (\Delta_a \Delta_b - \Delta_h)\}} \right\},$$

we get

$$\begin{aligned} \frac{d}{dt}V(X_t, Y_t, Z_t) &\leq -\alpha(\|X\|^2 + \|Y\|^2 + \|Z\|^2) + k\alpha(\|X\| + \|Y\| + \|Z\|) \\ &= -\frac{\alpha}{2}(\|X\|^2 + \|Y\|^2 + \|Z\|^2) - \frac{\alpha}{2} \{(\|X\| - k)^2 + (\|Y\| - k)^2 + (\|Z\| - k)^2\} + \frac{3\alpha}{2}k^2 \\ &\leq -\frac{\alpha}{2}(\|X\|^2 + \|Y\|^2 + \|Z\|^2) + \frac{3\alpha}{2}k^2, \quad \text{for some } k, \alpha > 0. \end{aligned}$$

This completes the proof. \square

Example 2 Now, as a special case of system (1.1) (for $P(t) \neq 0$), let us take $n = 2$ that $A, B, H(X(t - r(t)))$ defined in Example 1 hold. If we take $r(t) = \frac{1}{2146+t^2}$, then $0 \leq \frac{1}{2146+t^2} < \gamma$, and $r'(t) = \frac{-2t}{(2146+t^2)^2} \leq \xi$, $0 < \xi < 1$. Let

$$P(t) = \begin{pmatrix} \frac{1}{1+t^2} \\ \frac{1}{1+t^2} \end{pmatrix}, \quad \beta = \frac{1}{6} \quad \text{and} \quad \xi = \frac{1}{2},$$

we have that

$$\|P(t)\| = \frac{2}{1+t^2} \leq 2 \quad \text{and} \quad \gamma < \min \left\{ \frac{2}{13}, \frac{1}{6}, \frac{1}{2144} \right\}.$$

Thus, all the conditions of Theorem 4.2 are satisfied.

References

- [1] Afuwape, A. U.: *Ultimate boundedness results for a certain system of third-order nonlinear differential equations*. J. Math. Anal. Appl. **97** (1983), 140–150.
- [2] Afuwape, A. U., Omeike, M. O.: *Further ultimate boundedness of solutions of some system of third-order nonlinear ordinary differential equations*. Acta Univ. Palacki. Olomuc., Fac. rer. nat., Math. **43** (2004), 7–20.
- [3] Afuwape, A. U., Omeike, M. O.: *On the stability and boundedness of solutions of a kind of third order delay differential equations*. Applied Mathematics and Computation **200** (2000), 444–451.
- [4] Burton, T. A.: *Volterra Integral and Differential Equations*. Academic Press, New York, 1983.
- [5] Burton, T. A.: *Stability and Periodic Solutions of Ordinary and Functional Differential Equations*. Academic Press, Orlando, 1985.
- [6] Burton, T. A., Zhang, S.: *Unified boundedness, periodicity and stability in ordinary and functional differential equations*. Ann. Math. Pura Appl. **145** (1986), 129–158.
- [7] Ezeilo, J. O. C., Tejumola, H. O.: *Boundedness and periodicity of solutions of a certain system of third-order nonlinear differential equations*. Ann. Math. Pura Appl. **74** (1966), 283–316.
- [8] Ezeilo, J. O. C.: *n-dimensional extensions of boundedness and stability theorems for some third-order differential equations*. J. Math. Anal. Appl. **18** (1967), 395–416.
- [9] Ezeilo, J. O. C., Tejumola, H. O.: *Further results for a system of third-order ordinary differential equations*. Atti. Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. **58** (1975), 143–151.
- [10] Hale, J. K.: *Theory of Functional Differential Equations*. Springer Verlag, New York, 1977.
- [11] Sadek, A. I.: *Stability and Boundedness of a Kind of Third-Order Delay Differential System*. Applied Mathematics Letters **16** (2003), 657–662.
- [12] Sadek, A. I.: *On the stability of solutions of certain fourth order delay differential equations*. Applied Mathematics and Computation **148** (2004), 587–597.
- [13] Tiryaki, A.: *Boundedness and periodicity results for a certain system of third-order nonlinear differential equations*. Indian J. Pure Appl. Math. **30**, 4 (1999), 361–372.
- [14] Zhu, Y.: *On stability, boundedness and existence of periodic solution of a kind of third-order nonlinear delay differential system*. Ann. of Diff. Eqs. **8**, 2 (1992), 249–259.
- [15] Yoshizawa, T.: *Stability Theory by Liapunov's Second Method*. The Mathematical Society of Japan, Tokyo, 1996.